

Sample preparation checklist for the referring laboratory



Biomnis

- ► Patient must have fasted before samples collection.
- Tubes with phase separator (gel) are prohibited.
- It is essential to state the nature of the sample on each tube sent.

Category of tubes:



Li heparin







EDTA Dry Sodium citrate





Pot with stools

MiniCheck

Tubes to be collected



4 dry tubes

1 tube sodium citrate

Pretreatment of samples

Tubes to be sent (7 tubes)

- 2 x 2 mL refrigerated serum
- 2 x 1 mL refrigerated serum
- 2 x 2 mL frozen serum * <4h
- 1 mL frozen citrated plasma * <1h</p>

BasicCheck

Tubes to be collected

1 tube EDTA

5 dry tubes

1 tube sodium citrate

Pretreatment of samples

Tubes to be sent (7 tubes)

- 3 mL refrigerated EDTA whole blood
- 2 x 4 mL refrigerated serum
- 1 mL frozen serum ※ < 4h</p>
- 2 mL frozen serum * < 4h
- 1 mL frozen serum (protect from direct light) ※ < 1h30
- 1 mL frozen citrated plasma 🛠 < 1h

AgeCheck Woman

Tubes to be collected

4 dry tubes

1 tube sodium citrate

Pretreatment of samples

Tubes to be sent (4 tubes)

- 5 mL refrigerated serum.
- 2 x 2 mL frozen serum * < 4h</p>
- 1 mL frozen citrated plasma 🕸 < 1h

AgeCheck Man

Tubes to be collected

4 dry tubes

1 tube sodium citrate

Pretreatment of samples

Tubes to be sent (4 tubes)

- 4 mL refrigerated serum
- 2 x 2 mL frozen serum 🗱 < 4h
- 1 mL frozen citrated plasma ※ < 1h</p>

DermaCheck

Tubes to be collected

3 dry tubes

Urine (1st miction of morning)

Pretreatment of samples

Tubes to be sent (4 tubes)

- 5 mL refrigerated serum
- 1 mL frozen serum * < 4h</p>
- 1 mL frozen serum (protect from direct light) 🗱 < 1h30
- 4 mL of refrigerated urine

DigestCheck

To be collected

Stools

1 refrigerated stool sample

To be sent

FertiCheck Man

Tubes to be collected

1 tube sodium citrate

1 tube EDTA

4 dry tubes

Pretreatment of samples

Tubes to be sent (5 tubes)

- 3 mL refrigerated EDTA whole blood
- 4 mL refrigerated serum
- 2 mL frozen serum (protect from direct light) 🕸 < 1h30
- 🛡 2 mL frozen serum 🕸 < 4h
- 1 mL frozen citrated plasma 🗱 < 1h

For the unitary Juvenalis analyses, please prepare the sample for analysis as indicated on the Eurofins Biomnis test guide. For intestinal microbiota, you are required to order a special kit from our Eurofins Biomnis Connect client portal..



Sample preparation checklist for the referring laboratory



Biomnis

- ► Patient must have fasted before samples collection.
- Tubes with phase separator (gel) are prohibited.
- It is essential to state the nature of the sample on each tube sent.

Category of tubes:



Li heparin



Pretreatment

of samples









FertiCheck Woman

Tubes to be collected

- 2 tubes EDTA
- 4 dry tubes
- 1 tube sodium citrate

Tubes to be sent (7 tubes)

- 2 x 3 mL EDTA whole blood
- 2 x 3 mL refrigerated serum.
- 2 mL frozen serum (protect from direct light) * < 1h30</p>
- 2 mL frozen serum * < 4h</p>
- 1 mL frozen citrated plasma ※ < 1h
 </p>

Minerals

Tubes to be collected

- 2 dry tubes
- Urine (1st miction of morning)

Pretreatment of samples

Pretreatment

of samples

Tubes to be sent (2 tubes)

- 3 mL refrigerated serum.
- 4 mL refrigerated urine

NutriCheck

Tubes to be collected

- 1 tube EDTA
- 3 dry tubes
 - 1 tube sodium citrate

Tubes to be sent (5 tubes)

- 3 mL refrigerated EDTA whole blood
- 2 x 3 mL refrigerated serum
- 1 mL frozen serum (protect from direct light) 🗱 < 1h30
- 🕽 1 mL frozen citrated plasma 💥 < 1h

OxvCheck

Tubes to be collected

- 2 dry tubes
 - 3 tubes of Li heparin

Pretreatment of samples

Pretreatment

of samples

Tubes to sent (6 tubes)

- 1 mL refrigerated serum
- 1 mL frozen serum * <4h</p>
- 2x 2 mL refrigerated Li heparin plasma ● 2 mL frozen Li heparin plasma * <1h30</p>
- 1 pellet lithium heparin : frozen ※ <¹h</p>

VeggieCheck

Tubes to be collected

- 2 tubes EDTA
 - 3 dry tubes
- 1 tube sodium citrate
- Urine (1st miction of morning)

Tubes to be sent (6 tubes)

- 2 x 3 mL EDTA whole blood
- 4 mL refrigerated serum
- 1 mL frozen serum (protect from direct light) 🗱 < 1h30
- 1 mL frozen citrated plasma * < 1h</p>
- 4 mL refrigerated urine

Vitamins

Tubes to be collected

- 4 tubes EDTA
- 1 dry tube
- 1 tube of Li heparin

Tubes to be sent (7 tubes)

- 3 mL refrigerated EDTA whole blood
- 3 x 3 mL EDTA whole blood (protect from direct light) 🗱 < 4h
- 1 mL refrigerated serum.
- 1 mL frozen serum (protect from direct light) 🗱 < 1h30
- 🔍 2 mL frozen Li heparin plasma (protect from direct light) 🗱 < 4h

Nutritol™ food intolerances

Tubes to be collected

2 dry tubes

Pretreatment of samples

Pretreatment

of samples

Tubes to be sent (1 tube)

3 mL refrigerated serum